

COVID-19

Clinical Management of COVID-19 in Hospitalised Adults (including in pregnancy)

Introduction

Updated **6 May 2022** – Next planned update **24 June 2022**

- Revision of advice regarding deterioration ([page 3](#))
- Addition of advice to assess eligibility criteria for antivirals on hospital discharge ([page 3](#))
- Updated eligibility criteria for antivirals with addition of Down syndrome and sickle cell disease ([page 4](#))
- ****NEW**** figure added that provides a 'Heatmap' of eligibility for antivirals based on risk ([page 5](#))
- Access criteria and information for molnupiravir has been added ([page 6](#))
- Advice with pregnancy updated ([page 6](#))

New content in this update is highlighted in red.

This guideline is intended to be an accessible summary of hospital management of **ADULTS** (including in pregnancy) with **confirmed or probable COVID-19**. It has been adapted from international 'living' guidelines for the New Zealand context by an advisory group of New Zealand clinicians, including representation from Infectious Diseases, Respiratory Medicine, Intensive Care, General Medicine, Obstetric Medicine, Primary Care, Emergency Medicine and Pharmacy.

New evidence informing the optimal management of patients with COVID-19 continues to accumulate rapidly. This document will be reviewed and updated periodically, or in response to significant changes in evidence and/or recommendations by international guideline groups. Download the Ministry of Health [Āwhina app](#) to be notified when guideline updates are made.

Further detailed guidance, including treatments that are either not recommended, or recommended only within a clinical trial, can be accessed in links to international guidance at the end of this document. Local specialist services with expertise in managing COVID-19 may consider emerging treatments outside of this guideline.

Additional details or differences on managing a pregnant woman are highlighted in orange rows just below the adult guidance.

COVID-19

Lastly, the management of COVID-19 in [severely immunocompromised patients](#) presents unique challenges that are outside the scope of this guideline. Specialist advice from a patient's primary specialist **and** an Infectious Diseases physician is strongly recommended.

ASSESSMENT & DEFINITION		MILD		MODERATE		SEVERE / CRITICAL	
		SEVERITY CATEGORY	CLINICAL DEFINITION	SEVERITY CATEGORY	CLINICAL DEFINITION	SEVERITY CATEGORY	CLINICAL DEFINITION
CLINICAL DEFINITION		No symptoms	Any COVID-19 symptoms <u>without</u> features of pneumonitis	A clinically stable patient with any evidence of COVID-19 pneumonitis: ▪ New onset (or worsening) shortness of breath OR ▪ Infiltrates on plain chest radiograph OR ▪ Hypoxaemia that is EITHER... ...mild (92-94%), transient, or exercise-induced only (i.e. not requiring continuous oxygen therapy) OR ...sustained but able to maintain $\geq 92\%$ ($\geq 90\%$ for patients with chronic lung disease) with up to 4L/min oxygen via standard nasal prongs	Any of the following: ▪ Requiring CPAP or high-flow nasal oxygen to maintain saturations $\geq 92\%$ OR ▪ Acute respiratory distress e.g. RR > 30 OR ▪ Rapidly deteriorating clinical trajectory	Any of the following: ▪ Requiring mechanical ventilation to maintain saturations $\geq 92\%$ OR ▪ Requiring advanced circulatory support	
STAGE OF INFECTION		Almost all cases in the first 5 days; throughout in most vaccinated patients without risk factors.		Progression to moderate/severe disease most commonly develops ~5-7 days post onset of illness in patients with significant risk factors; the trajectory of deterioration can sometimes be rapid.			
SITE OF COVID CARE		Community		Individualised decision making	Hospital		
RESPIRATORY SUPPORT		Nil		Oxygen via NP	CPAP (or HFNO)	Mechanical ventilation	
VTE PROPHYLAXIS		Nil	Low dose enoxaparin <i>If hospitalised</i>	Low dose enoxaparin (or consider therapeutic-dose)	Low dose enoxaparin		
CORTICOSTEROIDS		Nil	Consider inhaled budesonide <i>If meets high risk criteria</i>	Dexamethasone	Dexamethasone		
ANTI-VIRAL THERAPY		Nil	Paxlovid® (or remdesivir) <i>If <5 days of illness AND meets high risk criteria</i>	Nil	Nil		
ANTIBODY THERAPY		Nil		Nil	Nil		
IMMUNE MODULATION		Nil		Baricitinib or tocilizumab	Tocilizumab		

Figure 1 - Severity assessment and therapeutic options in COVID-19 management.

Initial management

	MILD	MODERATE	SEVERE / CRITICAL
DEFINITION	No symptoms OR URTI symptoms only OR cough, new myalgia or asthenia <u>without</u> new shortness of breath or reduction in resting oxygen saturation	Stable adult patient presenting with shortness of breath and/or reduction in resting oxygen saturation while breathing air. Able to maintain oxygen saturation $\geq 92\%$ (or $\geq 90\%$ for patients with chronic lung disease) with up to 4 L/min oxygen via nasal prongs	Adult patients meeting any of the following criteria: • Respiratory rate ≥ 30 /min • Oxygen saturation $< 92\%$ on 4L/min oxygen via nasal prongs • Clinically deteriorating
Pregnancy: use an oxygen saturation target of $\geq 94\%$ rather than $\geq 92\%$			
BASELINE TESTING & WORK-UP	<ul style="list-style-type: none"> Pulse oximetry Other tests only as clinically indicated Low value testing is discouraged 	<ul style="list-style-type: none"> FBC, Creat, electrolytes, LFTs, CRP ECG only if specific indication Chest x-ray Venous blood gas (consider arterial) Investigations for CAP (e.g. urinary antigens, sputum PCR) 	<ul style="list-style-type: none"> FBC, Creat, electrolytes, LFTs, CRP ECG Chest x-ray Venous blood gas (consider arterial) Investigations for CAP (e.g. urinary antigens, sputum PCR)

COVID-19

		<p>panel) if radiography suggests bacterial infection</p> <ul style="list-style-type: none"> Consider d-dimer & ferritin 	<p>panel) if radiography suggests bacterial infection</p> <ul style="list-style-type: none"> Blood cultures if febrile or shocked Coag screen, d-dimer, ferritin, BNP, Troponin
	<ul style="list-style-type: none"> Note – in vaccinated individuals with Omicron variant infection, COVID-19 may not be the primary diagnosis responsible for hospital presentation. It is important to consider concurrent non-COVID-19 medical conditions during evaluation. 		
	<ul style="list-style-type: none"> Pregnancy: also request urine protein:creatinine ratio, coagulation profile, group and screen (or cross match if delivery is thought to be imminent) <i>NB</i> CXR and CT chest / CTPA can safely be performed in pregnancy if clinically indicated. Laboratory results should be cautiously interpreted, using pregnancy-specific ranges where available. There are no validated pregnancy-specific values for D-dimer; consider monitoring trend for prognostic purposes in severe/deteriorating COVID-19. 		
TREATMENT ESCALATION PLANNING	<ul style="list-style-type: none"> Assess ability to safely isolate in community. Notify and refer through local pathways Consider & document risk factors for severe COVID-19 	<ul style="list-style-type: none"> Assess & document individual risk factors for poor outcome Early discussion of patient goals of care, including existing advanced care plans, with patient and their family/whānau Early, clear documentation of resuscitation decision and treatment escalation plan for <u>all</u> patients, specifically including appropriate modalities of respiratory support 	
	<ul style="list-style-type: none"> NOTE – any new deterioration > 5 days post onset of illness requires careful assessment. Deterioration to severe COVID-19 can occur rapidly. Pneumonitis continues to develop in the second (or sometimes third) week of illness, particularly in older or unvaccinated patients 		
	<ul style="list-style-type: none"> For pregnant and post-partum observations, utilise a maternity-specific chart (if available) Recommend early consultation with Obstetrics, Anaesthesia and NICU (and Obstetric Physician if available) 		
DISPOSITION DECISION	<ul style="list-style-type: none"> Encourage discharge Offer COVID-19 treatment on discharge if meet eligibility criteria Liaise with local Public Health Unit or Regional Isolation and Quarantine (RIQ) according to regional processes 	<ul style="list-style-type: none"> Discuss with local COVID team Admit to hospital if SaO₂ <93% Consider discharge if SaO₂ ≥93% according to local protocols and availability of acute community COVID-19 care (e.g. primary care or hospital in the home service) Offer COVID-19 treatment on discharge if meet eligibility criteria 	<ul style="list-style-type: none"> Admit to hospital ICU and/or Respiratory review
MONITORING & MARKERS OF CLINICAL DETERIORATION	<ul style="list-style-type: none"> Risk of deterioration is significantly reduced by vaccination and infection with Omicron variant. Individualised risk assessment should include consideration of vaccination status, day of illness, age, immunocompromise and comorbidities that increase risk of severe disease. Ferritin and d-dimer are suggested as severity/prognosis markers, as part of an overall assessment Monitor for progressive respiratory failure and sepsis, especially after day 5 of illness Only repeat CXR during admission for confirmed COVID-19 for specific clinical indications Perform a chest CT scan only if it would change management, in particular if concern for pulmonary embolism Anticipate complications such as delirium, pulmonary embolism, other thromboembolism, arrhythmias, cardiac impairment, acute kidney injury, sepsis, shock and multi-organ dysfunction, and address using existing standards of care. Also be aware of potential medication complications Repeat baseline investigations periodically in patients who are not clearly improving, in order to detect & manage the above complications 		

	<p>Additional considerations in pregnancy:</p> <ul style="list-style-type: none"> • Screen for pre-eclampsia in all pregnancies > 20/40 gestation and review at each assessment: i.e., systolic BP \geq 140mmHg and/or diastolic \geq 90, worsening peripheral oedema, headache, visual changes or upper abdominal pain. Risk of pre-eclampsia is increased in COVID-19. • Consider repeating laboratory investigations if there is a change in maternal condition • Appropriateness and frequency of foetal heart rate monitoring and ultrasound to be considered on an individual basis, accounting for gestational age and clinical severity. Consider parameters for delivery (in discussion with neonatologists, anaesthetists \pm intensive care team) • Consider steroids for the fetal lung maturation, and magnesium sulphate for neuroprotection or severe pre-eclampsia as per local obstetric guidelines • Consider emergency delivery if required for maternal resuscitation (including need for prone positioning) or for immediate foetal concern
NOTIFICATION	<ul style="list-style-type: none"> • Discuss all admitted cases with local COVID team at the earliest opportunity, according to local protocols • If not already notified according to regional process, (e.g. by laboratory) contact local Public Health Unit <p>If hospitalised, all pregnant women should have multidisciplinary assessment by obstetricians, midwives, neonatologists +/- an obstetric physician at the earliest opportunity</p> <ul style="list-style-type: none"> • Recommend notification of all antenatal and postnatal cases to New Zealand Registry of COVID-19 in Pregnancy
CLINICAL TRIALS	<ul style="list-style-type: none"> • As the optimal management of COVID-19 is not yet known, the standard of care is to be offered enrolment in a clinical trial, if available • All patients should be screened for eligibility for a locally available COVID-19 clinical trial (e.g. 'REMAP-CAP' and 'ASCOT-ADAPT')

COVID-19 Therapeutics: patients **not requiring oxygen**

There are increasing therapeutic options for COVID-19 in patients who do not require oxygen. The main benefit of these treatments is in reducing hospitalisation **with severe COVID-19**, with a possible small reduction in mortality. These treatments have not been studied in vaccinated individuals or in people infected with the Omicron variant, both of which are anticipated to significantly reduce the benefit of treatment compared with unvaccinated study cohorts. When stocks are limited, it is therefore of critical importance that treatments are prioritised to patients with the highest absolute risk of severe COVID-19. The Pharmac [recently released access criteria](#) for antiviral treatments outline groups who are at high absolute risk of hospitalisation in New Zealand. As such, **we recommend that all treatments** (including antivirals and budesonide) **for patients not requiring oxygen be prioritised to people who:**

- 1) **Have** at least **five of the following risk factors** (*summarised in 'heat map' below*)
 - a. Any combination of the **risk factors for severe COVID-19 disease**¹ (with each individual condition counting as one risk factor)
 - b. Māori or any Pacific ethnicity (**counts as one risk factor**)
 - c. Patient is aged 65 years and over (**counts as two risk factors, or three if has not completed a full course of vaccination**) OR is 50 years and over and has
 - d. Not completed a full course of vaccination² (**counts as one risk factor**)
- 2) OR are **severely immunocompromised**³ and not expected to reliably mount an adequate immune response to COVID-19 vaccination or SARS-CoV-2 infection, regardless of vaccination status
- 3) OR have either **Down syndrome OR sickle cell disease**

COVID-19

For these treatments, patients should not already have COVID-19 associated pneumonitis requiring oxygen. If a patient requires oxygen for COVID-19, **different therapeutics recommendations** apply.




Figure 2: Heatmap of eligibility for antivirals based on risk

Notes:

¹ **Risk factors** are detailed on the Ministry of Health (MOH) website and include: obesity, chronic lung disease, chronic kidney disease, heart disease, diabetes, hypertension, chronic liver disease, active malignancy, chronic neurologic disease and severe mental health illness.

² **Incomplete vaccination** is defined as fewer than two doses by the linked [Ministry of Health document](#). However, for the purposes of this guideline, we currently consider incomplete vaccination to be:

- Fewer than 2 doses of vaccine
- OR 2 doses of vaccine, with second dose < 7days or > 6months before symptom onset

³ The definition of **immunocompromise** in PHARMAC access criteria aligns with the eligible population for a **three-dose primary vaccine** series. However, a subgroup of **severely immunocompromised** individuals are at higher risk of severe outcomes, including:

- Solid organ transplant recipient, particularly if within 12 months of transplantation, if requiring more than routine maintenance immunosuppression, treated with mycophenolate mofetil, or treated for rejection within past 12 months
- Within 24 months of haematopoietic stem cell transplant or CAR-T cell therapy.
- Graft-versus-host disease treated with multi-modal immunosuppressive therapy
- Treated B-cell haematologic malignancy (e.g. multiple myeloma, chronic lymphocytic leukaemia, lymphoma) within the past 6 months
- Receipt of anti-CD20 monoclonal antibody therapy (e.g. rituximab) within the past 12 months
- Primary or acquired hypogammaglobulinaemia (IgG <3), even if now on replacement immunoglobulin

COVID-19

- Primary immunodeficiency associated with severe B-cell or combined cellular defects
- Advanced HIV with CD4 <200
- Other conditions (on case by case basis) felt to have profound immunocompromise on the basis of combined immunosuppression, functionally equivalent to the above groups

MODALITY	PATIENT SUB-GROUPS	RECOMMENDATION
ANTIVIRALS	Adults who meet high risk criteria <u>AND</u> are within 5 days of symptom onset <u>AND</u> do not have severe hepatic (Childs-Pugh class C) or renal impairment (eGFR <30ml/min) <u>AND</u> do not have a potentially serious drug-drug interaction with ritonavir	Give Paxlovid (nirmatrelvir and ritonavir):* <ul style="list-style-type: none"> • (nirmatrelvir 300mg + ritonavir 100mg) PO q12h for 5 days • eGFR 30-59ml/min: nirmatrelvir 150mg + ritonavir 100mg po q12h for 5 days • Use barrier contraception for 7 days after last dose <p>*NB Paxlovid prescriber advice available here. Management of common drug interactions highlighted here.</p>
	Adults who meet high risk criteria <u>AND</u> are unable to receive Paxlovid <u>AND</u> are within 7 days of symptom onset <i>Guidance for further prioritisation of remdesivir to patients at highest risk is available here.</i>	Consider remdesivir: <ul style="list-style-type: none"> • 200mg IV on day 1, then 100mg IV q24h for further 2 days (maximum 3 days total) • Limited data of safety in patients with eGFR <30ml/min or peritoneal dialysis.* Use if benefits felt to clearly outweigh potential risks. Likely to be safe in haemodialysis. <p>*Consider a two dose regimen (i.e. omission of day 3 dose) for patients with eGFR<30: modelling suggests this may provide equivalent drug concentrations to patients with normal renal function.</p>
	Adults who meet high risk criteria <u>AND</u> are unable to receive Paxlovid <u>AND</u> are unable to receive remdesivir <u>AND</u> are within 5 days of symptom onset	Consider molnupiravir #: <ul style="list-style-type: none"> • 800mg PO q12h for 5 days • Use barrier contraception while taking molnupiravir and for 4 days after last dose <p>#NB molnupiravir is less effective at reducing risk of severe COVID-19 than other antivirals, and is not recommended for regular use in high-risk patients presenting to hospital.</p>
	Adults with COVID-19 after day 7 of illness	<ul style="list-style-type: none"> • Do not start antivirals • Complete course if started earlier in illness
	Recommend discuss all severely immunocompromised patients with Infectious Diseases	
Pregnancy (meeting the same clinical criteria as above)	<ul style="list-style-type: none"> • Do not use Paxlovid or molnupiravir in pregnancy • Avoid breastfeeding during and for 7 days after Paxlovid or 4 days after molnupiravir • Use remdesivir if >12/40 gestation as per Adult Guideline indications with the same dosing as above • Remdesivir is compatible with breastfeeding 	
STEROIDS	Adults who meet high risk criteria <u>AND</u> are ineligible for antivirals <u>AND</u> are within 14 days of symptom onset	Consider inhaled budesonide 800micrograms BD for up to 14 days if respiratory symptoms <ul style="list-style-type: none"> • Updated budesonide guidance available here <p>Do not use systemic steroids to treat COVID-19 without an oxygen requirement</p>

	Adults without oxygen requirement, but another evidence-based indication for systemic steroids (e.g. asthma/COPD exacerbations)	Steroids as per usual practise
ANTIBODY THERAPY	Adults with any severity of illness	Do not use casivirimab/imdevimab (Ronapreve) due to lack of efficacy against Omicron, and negligible Delta variant transmission in New Zealand currently *Guidance about use in specific cases if advised by an expert clinician is available here .

Supportive management: **all patients in hospital**

MODALITY	PATIENT SUB-GROUPS	RECOMMENDATION
RESPIRATORY SUPPORT	All patients	<ul style="list-style-type: none"> • Switch nebulisers to metered dose inhalers via spacer if possible • Monitor closely for worsening hypoxia if elevated work of breathing or respiratory rate
	SpO ₂ <92% at rest	<ul style="list-style-type: none"> • Administer dry oxygen (1-4 L/min) via standard nasal prongs • Use Hudson mask (5-10 L/min), Venturi device or high flow nasal oxygen (HFNO) if required • Aim for SpO₂ 92–96% (88–92% for those at risk of hypercapnic respiratory failure) • Encourage use of self-proning
	Unable to maintain SpO ₂ ≥92% on conventional oxygen or high flow nasal oxygen [HFNO] (requiring FiO ₂ >40%)	<ul style="list-style-type: none"> • Consider CPAP. Settings should be individualised, but a starting pressure of 8-10cm H₂O is common • Continue HFNO if CPAP unavailable, during meal breaks from CPAP or patient intolerance of CPAP • Encourage use of self-proning
	Hypercapnic patients with underlying COPD or OHS	<ul style="list-style-type: none"> • Consider BiLevel Non-Invasive Ventilation (NIV) in addition to above
	Pregnancy	<ul style="list-style-type: none"> • SpO₂ target is ≥ 94%; ideally aim for 96-98%. • After 20/40 avoid positioning flat on back: use a wedge for lateral supine positioning. Left lateral during resuscitation or if hypotensive. • Self-proning may be possible (depending on gestation and habitus).
FLUID MANAGEMENT	<ul style="list-style-type: none"> • Assess for hypovolaemia and correct as required. • Avoid excessive resuscitation or 'maintenance' fluids • Anticipate and monitor ongoing fluid losses 	
VTE PROPHYLAXIS	<ul style="list-style-type: none"> • All patients enrolled in ASCOT-ADAPT or REMAP-CAP (anticoagulation domains) 	<ul style="list-style-type: none"> • As per trial protocol
	Hospitalised adults with: <ul style="list-style-type: none"> • mild COVID-19 • <u>OR</u> severe and critical COVID-19 <p><u>AND</u> no contra-indication to anticoagulation e.g. risk for major bleeding</p>	Enoxaparin 40mg SC once daily (standard prophylaxis) <ul style="list-style-type: none"> • Adjust dose for impaired renal function <p>(NB Therapeutic-dose anticoagulation is not beneficial and probably hazardous when initiated prophylactically in severe and critical COVID-19)</p>

COVID-19

	<p>Hospitalised adults with moderate COVID-19 AND no contra-indication to anticoagulation e.g. risk for major bleeding</p> <p>(NB moderate = <i>stable adult patient presenting with shortness of breath and/or reduction in resting oxygen saturation while breathing air. Able to maintain oxygen saturation $\geq 92\%$ (or $\geq 90\%$ for patients with chronic lung disease) with up to 4 L/min oxygen via nasal prongs</i>)</p>	<p>Therapeutic dose anticoagulation should be considered over standard prophylaxis for up to 14 days, or until clinical recovery (discharge or resolved hypoxia)</p> <p>Enoxaparin 1mg/kg SC twice daily (max 150mg BD)</p> <ul style="list-style-type: none"> Adjust dose for impaired renal function <p>All other patients should receive standard prophylaxis as detailed above</p>
	<p>Hospitalised pregnant adults with mild OR severe/critical COVID-19 UNLESS:</p> <ul style="list-style-type: none"> Delivery expected within 24 hours (if only on enoxaparin 40mg SC once daily then 12 hourly) Platelets < 50 Actively bleeding / coagulopathy Severe hypertension (>160/110) 	<p>Enoxaparin 40mg SC once daily (standard prophylaxis)</p> <ul style="list-style-type: none"> dose adjustment may be necessary if current weight $\geq 90\text{kg}$
	<p>Hospitalised pregnant or postpartum adults with moderate COVID-19 AND no contra-indication to anticoagulation (as above)</p>	<p>Consider therapeutic anticoagulation as for non-pregnant adults (above)</p>
<p>Anticoagulation in pregnancy should be considered for a longer duration if post-partum or has additional risk factors for VTE (discuss with Obstetrics)</p>		
<p>INTENSIVE CARE</p>	<p>Regular, open and early discussions between ward-based clinicians and local ICU team is strongly encouraged. In addition to local referral guidelines, ICU review should be prompted by the following:</p> <ul style="list-style-type: none"> Significant oxygen requirement (e.g. requiring FiO₂ of >40% to maintain SpO₂ >92%, or needing CPAP) Increased work of breathing with impending respiratory failure Haemodynamically unstable and / or hypotension not responsive to fluid bolus Rapidly worsening tachypnoea or hypoxaemia <p>Detailed clinical guidance for ICU care of COVID-19 is beyond the scope of this guideline</p>	
<p>ANTIBIOTIC THERAPY</p>	<p>Antibiotics should not be used to treat COVID-19 pneumonitis: bacterial co-infection is uncommon.</p> <p>Severe/critical COVID-19 especially with any deterioration occurring >7 days post onset and/or >3 days after hospital admission</p>	<ul style="list-style-type: none"> Evaluate for secondary infection, including hospital-acquired infection Discuss with local Infectious Diseases / Microbiology team if concern for secondary infection
<p>COMMUNICATION & HOLISTIC CARE</p>	<p>Encourage for all patients:</p> <ul style="list-style-type: none"> Clearly communicate typical symptoms and anticipated clinical course with patient, family/whānau or carers Reinforce importance of complying with all Public Health messages, including self-isolation and testing When possible, explain risks, benefits and likely outcomes of treatments with patients, family/whānau or carers Use an interpreting service to assist communication if required Facilitate regular clinical updates, and video calls between patient family/whānau or carers Routinely refer to local cultural and/or spiritual support services 	

COVID-19

	<ul style="list-style-type: none"> • Consider early involvement of Palliative Care and/ or Liaison Psychiatry services to assist with symptom management, particularly anxiety, dyspnoea and delirium/agitation • Ensure appropriate housing, financial and social support is in place prior to discharge (including a working phone). If concerns, refer to social work • If welfare or cultural support issues identified, liaise with Public Health or Regional Isolation Quarantine (RIQ) according to regional processes as part of discharge planning 	
	<ul style="list-style-type: none"> • Ensure Maternity services including lead maternity carer are alerted so wrap-around antenatal and post-natal care can be provided for the mother and baby 	
THERAPIES FOR EXISTING INDICATIONS	<ul style="list-style-type: none"> • Nocturnal CPAP for Obstructive Sleep Apnoea (inpatients) 	Consider changing usual vented CPAP mask to a non-vented mask + expiratory port + filter (decision depends on equipment availability and staff expertise)
	<ul style="list-style-type: none"> • ACE-inhibitors / ARBs • Oral contraceptive pill (with or without oestrogen) 	<ul style="list-style-type: none"> • Usual care (i.e. may be continued in COVID-19 unless otherwise contra-indicated)
	<ul style="list-style-type: none"> • Corticosteroids for asthma/COPD (inhaled or oral, with or without bronchodilators) 	<ul style="list-style-type: none"> • Usual care • Do not use a nebuliser unless definite clinical need
	<ul style="list-style-type: none"> • Oral menopausal hormone therapy / HRT 	<ul style="list-style-type: none"> • Consider stopping until after recovery
	<ul style="list-style-type: none"> • All pregnancy-related supplements and medications should be continued 	
SURGERY	<ul style="list-style-type: none"> • Elective minor surgery should generally be deferred until at least four weeks, and major surgery until 8-12 weeks, following recovery from COVID-19 if patient outcome is not compromised • Acute surgical procedures that are indicated during active COVID-19 infection should be discussed with local ID and infection control services • Comprehensive preoperative and ongoing assessment must be carried out to ensure optimal clinical decision-making 	
	<ul style="list-style-type: none"> • Caesarean section (including emergency) should not be deferred if clinically indicated, e.g. if needed for maternal resuscitation or immediate fetal concern; mode of delivery should otherwise remain based on obstetric indication 	

COVID-19 Therapeutics: patients **requiring oxygen**

MODALITY	PATIENT SUB-GROUPS	RECOMMENDATION
STEROIDS	Adults with sustained oxygen requirement	<p>Dexamethasone 6mg* daily PO/IV for 10 days OR until hospital discharge</p> <p>Do not routinely continue after discharge if completed at least 5 days in hospital</p> <p>*consider dexamethasone 12mg PO/IV on day 1 if would qualify for immunomodulation, but medication is unavailable within next 24 hours</p>

	<p>Pregnancy with sustained oxygen requirement to maintain SpO2 ≥94%</p>	<p>If steroids needed for fetal lung maturation (usually < 34⁺⁶ weeks):</p> <ul style="list-style-type: none"> dexamethasone 6mg IM every 12 hours for four doses THEN either prednisone 40mg PO daily, OR hydrocortisone 80mg IV twice daily <p>If steroids <u>not</u> required for fetal lung maturation, use non-fluorinated steroids:</p> <ul style="list-style-type: none"> prednisone 40mg PO daily OR hydrocortisone 80mg IV twice daily OR methylprednisolone 40mg IV once daily <p>Total duration is 10 days total OR until discharge, whichever is sooner.</p>
<p>Risk of gestational diabetes: monitor blood glucose levels closely and start treatment if elevated.</p>		
<p>ANTIVIRAL THERAPY</p>	<p>All patients enrolled in ASCOT-ADAPT trial (anti-viral domain)</p>	<p>As per trial protocol & randomisation (in addition to remdesivir, if indicated below)</p>
	<p>Adults with new sustained oxygen requirement within first 7 days of illness</p>	<ul style="list-style-type: none"> Do not start remdesivir Complete course (3 days) if started earlier in illness
	<p>Adults with COVID-19 after day 7 of illness</p>	<ul style="list-style-type: none"> Do not start remdesivir Complete course (3 days) if started earlier in illness
	<p>Adults with severe immunocompromise with any stage/severity of COVID-19</p>	<p>Discuss with local infectious diseases team</p>
	<p>Pregnancy (meeting the same clinical criteria as above)</p>	<ul style="list-style-type: none"> Use remdesivir if >12/40 gestation as per Adult Guideline indications with the same dosing as above Remdesivir is compatible with breastfeeding
<p>ANTIBODY THERAPY</p>	<p>Adults with any severity of illness</p>	<p>Do not use casirivimab/imdevimab (Ronapreve) due to lack of efficacy against Omicron, and negligible Delta variant transmission in New Zealand currently</p> <p>*Guidance about use in specific cases if advised by an expert clinician is available here.</p>
<p>IMMUNE MODULATION THERAPY</p>	<p>In patients receiving systemic steroids in combination with immune modulation, we recommend screening for, and consider empiric treatment of latent infection, e.g. Hepatitis B or strongyloidiasis (in patients who have lived in an endemic region)</p>	
	<p>There are no trials of immune modulation therapies currently recruiting in New Zealand</p> <p>Adults with moderate COVID-19</p> <ul style="list-style-type: none"> AND receiving systemic steroids AND elevated CRP or other evidence of severe systemic inflammation OR clinically deteriorating AND there is not another active, severe concurrent infection 	<p>Give baricitinib:</p> <ul style="list-style-type: none"> 4mg PO/NG daily for 14 days or until hospital discharge Reduce to 2mg PO daily if eGFR 30-60mL/min Reduce to 1mg PO daily if eGFR 15-29mL/min* Do not use if eGFR <15mL/min Avoid in pregnancy or breastfeeding Baricitinib is a section 29 product <p>OR tocilizumab:</p> <ul style="list-style-type: none"> 8mg/kg IV (actual body weight) rounded to nearest 80mg or 200mg vial (max dose 800mg), as a single dose Notes: risk of secondary infection may be increased; CRP response is inhibited, and consequent reduction in CRP does not necessarily correlate with response to treatment <p>*baricitinib 2mg PO every 48 hours is an alternative</p>

	Adults with severe / critical COVID-19 requiring non-invasive or mechanical ventilation or organ support: <ul style="list-style-type: none"> • <u>AND</u> receiving systemic steroids • <u>AND</u> there is not another active, severe secondary infection 	<p>Give tocilizumab as above</p> <ul style="list-style-type: none"> • Start as soon as possible if requiring NIV, mechanical ventilation or other organ support <p>OR baricitinib, if tocilizumab is unavailable (as above).</p> <ul style="list-style-type: none"> • If tocilizumab is available and baricitinib commenced earlier in illness, suggest change to tocilizumab <p>Do not treat with both baricitinib and tocilizumab together</p>
	COVID-19 not meeting the criteria above	Do not use immune modulation therapy
	Pregnancy (meeting the same clinical criteria as above)	<p>Give tocilizumab (same dosing as above):</p> <ul style="list-style-type: none"> • <i>Notes: Tocilizumab crosses the placenta after 28/40, but no evidence of harm to date. Suggest deferring live vaccination (i.e. Rotarix, BCG) up to 6 months in neonates with antenatal exposure. All other vaccinations are safe.</i> • <i>Compatible with breastfeeding.</i> • <i>May cause raised ALT and thrombocytopenia, mimicking pre-eclampsia / HELLP.</i> • <i>Do not use baricitinib (as above)</i>

Discharge Planning and Follow-up

Patients with Suspected, Probable or Confirmed COVID-19 who are being considered for discharge need to have specific decisions made about the following aspects of post-discharge care:

FURTHER INVESTIGATIONS	<ul style="list-style-type: none"> • Follow-up investigations are not universally required after COVID-19 • A repeat chest x-ray in 6-12 weeks to confirm resolution of pulmonary opacities should be arranged for individuals with significant radiographic abnormalities and / or risk factors for lung cancer
DISCHARGE DESTINATION	<ul style="list-style-type: none"> • Anyone with COVID symptoms or suspected infection being discharged before PCR results are available should be tested using a RAT. If the RAT (or a PCR) is negative, the person should be advised to stay home until symptoms resolve and seek a further test if symptoms worsen. • Anyone who tests positive (on RAT or PCR) should be able to be discharged home but should be linked in through the Care in the Community Model if needed for follow-up, and household members should be advised to self-isolate as per standard advice for positive cases. Note that positive RAT results need to be recorded in My Covid Record. • The local Medical Officer of Health does not need to be notified of discharge of a positive case.
CLEARANCE FROM ISOLATION	<ul style="list-style-type: none"> • The decision to end isolation should be consistent with Public Health policies and local hospital infection prevention and control policies, which may be different. • Local hospital isolation policy should be followed until point of discharge • Release from isolation after discharge should align with the current Public Health Policy for community isolation: this is now taken as 7 days from date of onset of symptoms or date of positive test (whichever is earlier, starting from day zero). • Exceptions to this duration may include severe immunocompromise and severe/critical COVID-19. It is advisable to seek the advice of an infectious disease specialist or microbiologist for severely immunocompromised individuals. Additional testing may be useful, such as serial NAAT/PCR testing suggestive of low viral load (i.e. negative or with high cycle threshold), high or increasing antibody levels or repeatedly negative RAT tests.
	<p>All patients</p> <p>Encourage vaccination if not completed eligible vaccination course (including booster).</p>

		<ul style="list-style-type: none"> • If not completed primary vaccination series before infection, vaccination is recommended from 4 weeks after clinical recovery, even if treated with anti-SARS-CoV-2 antibody therapy (convalescent plasma or monoclonal antibody such as Ronapreve) • If completed primary vaccination series before infection, booster vaccination is recommended from 12 weeks after clinical recovery <p>Educate about anticipated gradual recovery from COVID-19, and potential for persistent symptoms.</p> <ul style="list-style-type: none"> • Encourage those with persistent symptoms after 6 weeks to arrange assessment by their GP.
	Patients with significant respiratory failure (and/or persistent dyspnoea), or other persistent organ dysfunction	Specialist clinic follow-up, investigations and support following discharge (as advised by local specialty services)
	Pregnancy (or recently post-partum)	<ul style="list-style-type: none"> • VTE prophylaxis - refer to specific guidelines above • Recommend follow up growth scan within 2 weeks • If possible, delay follow-up CXR until post-partum

Links to other guidelines

- Australian COVID-19 living guidelines: <https://covid19evidence.net.au/>
- NICE (UK) living guideline: <https://www.nice.org.uk/guidance/ng191>
- National Institute of Health (USA): <https://www.covid19treatmentguidelines.nih.gov/>
- WHO COVID-19 living guideline: <https://www.who.int/publications/i/item/WHO-2019-nCoV-therapeutics-2022.1>
- Ontario COVID-19 Science Advisory Group guideline (Canada): <https://covid19-sciencetable.ca/sciencebrief/clinical-practice-guideline-summary-recommended-drugs-and-biologics-in-adult-patients-with-covid-19-version-10-0/>